

HUMIPLX[®]

GENERAL K BOOSTER

0-0-15



**CONTAINS (THA)
TECHNICAL HUMIC ACIDS**

GUARANTEED ANALYSIS

Soluble Potash (K₂O)..... 15.00%

Derived From: Potassium Thiosulfate.

Also contains NON-PLANT FOOD
0.8% Humic Acid derived from Leonardite.

KEEP OUT OF REACH OF CHILDREN

WARRANTY: WESTERN NUTRIENTS CORPORATION makes no warranty, express or implied, including the warranties of merchantability and/or fitness for any particular purpose, concerning this material, except those which are contained on the Western Nutrients Corporation label attached to the product container.

NET CONTENTS 5 GALLONS
18.93 LITERS
11 LBS. PER GAL @ 68° F
1108 GRAMS PER LITER @ 20° C



MANUFACTURED BY - WESTERN NUTRIENTS CORPORATION

245 Industrial Street, Bakersfield California 93307 • (661) 327-9604
(661) 327-1740 Fax • (800) 542-6664 Ca. Only / E-mail: westernnutrients@lightspeed.net
Website: <http://www.westernnutrientscorp.com>

PRODUCT INFORMATION

CROPS

HUMIPLX GENERAL BOOSTER Liquid Nutrients can be applied to most vegetable crops, row crops, deciduous fruit and nut trees, citrus, avocados, grapes, melons, ornamentals, turf, pasture, range grasses, and most other crops.

HUMIPLX GENERAL BOOSTER is a new liquid plant food developed for use as a foliar feed, a regular plant food applied to the soil, and as a starter plant food with the seed or transplant.

HUMIPLX GENERAL BOOSTER liquid nutrients contain ENHANCE[®] (THA) Technical Humic acids. HUMIPLX nutrients are beneficial in combination with plant food and nonphytotoxic when used as directed.

HUMIPLX nutrients with (THA) Technical Humic Acids are unique as they can be used in most all forms of liquid fertilizers. HUMIPLX nutrients can be banded at planting time, side-dressed or sprayed in water solutions directly on deficient plants. Humic Acids may aid in the uptake of micro-nutrients.

HUMIPLX liquid nutrients have corrected deficiencies of a great many row crops, vegetables, and ornamental plants under soil conditions ranging from high organic matter to very low organic matter and from strong acid soils (pH 3) to high alkaline soils (pH 8.5) containing considerable calcium carbonate (free lime). HUMIPLX can be effective under dry land and irrigation farming conditions. Differences in soil conditions, climate and plant varieties will determine how much more effective HUMIPLX liquid nutrients are than other sources of nutrients.

NOTE: These humic acid products are used to **fortify, produce, set and hold**. It is important that timely applications are made to achieve these results.

Nuts: 2-6 quarts per acre four times each season starting at first full leaf, then at fruit set, fruit size and fruit color.

Walnuts - Almonds: 2-6 quarts per acre three times each season starting at first full leaf.

Grapes: 2-6 quarts per acre five times each season starting at first full leaf, then past bloom, early bunch set, berry size and at berry color and just before harvest.

Cotton: 2-6 quarts per acre four times each season starting at first true leaf then first boll set, maximum boll set and two weeks before defoliation.

Tomatoes: 2-6 quarts per acre four times each season starting at first true leaf then at early fruit size, fruit color and one week before harvest.

Sugar Beets: 2-6 quarts per acre three times each season starting at first full leaf. Last application should be mid-season.

Alfalfa: 2-6 quarts per acre immediately after each cutting during the season and three to four gallons/acre after the final cutting of the season.

Wheat - Barley - Oats - Rice: 2-6 quarts per acre tillering and again at early cough stage.

Beans - Peas: 2-6 quarts per acre.

Onions - Garlic: 2-6 quarts per acre.

Corn: 2-6 quarts per acre three times each season at three week intervals.

Lettuce - Celery - Cole Crops: 2-6 quarts per acre four times each season starting at second true leaf and with the last application two weeks before harvest.

Asparagus: 2-6 quarts per acre two times while fern is full and two to four gallons/acre two weeks before fern turns yellow in the fall.

Strawberries: 2-6 quarts per acre at first early fruit set and after each picking.

Potatoes: 2-6 quarts per acre four to six times each season with the first at early emergence, then evenly spaced with the last application two weeks before leaf die-down.

Citrus: 2-6 quarts per acre three times year starting at early fruit set, then fruit size and early color.

Peppers, Cucumbers, Melons, and Squash: 2-6 quarts per acre Apply at 1-2 week intervals.

Transplant Solutions:

A. Mix one gallon in 100 gallons of water and drench roots (for vegetables drench the entire plant). Plant immediately after drenching. Do not allow plants to dry out or wilt.

Aircraft and low volume sprayers: Use a minimum of 10 gallons of water per acre.

Conventional sprayers: Use a minimum of 20 gallons of water per acre.

MINIMUM DILUTION 1 QUART PER 10 GALLONS OF WATER

HUMIPLX is a Registered Trademark of Western Nutrients Corporation.

ENHANCE is a Registered Trademark of Western Nutrients Corporation consisting of (THA) Technical Humic Acids.